Post Vaccination Advice Sheet for Parents

Diphtheria, tetanus, pertussis, polio, hib and	Rotavirus
 hep B (6 in 1) Pain, redness and swelling at the injection site. Fever. Vomiting. Abnormal crying. Irritability. Loss of appetite. 	 Crying Mild irritability. Fever. Diarrhoea. Vomiting.
Diphtheria, tetanus, pertussis, polio (4 in 1)	Pneumococcal (13 serotypes)
 Pain, redness and swelling at the injection site. Mild fever. Irritability. Loss of appetite. Restless. Increased crying. 	 Pain, redness and swelling at the injection site. Fever. Muscle, joint aches or pain. Chills. Headache. Nausea. Vomiting. Stiffness of the leg at site of vaccination.
Haemophilus Influenza type B (Hib) and Meningitis C (Men C)	Measles, Mumps and Rubella (MMR)
 Pain, redness and swelling at the injection site. Fever. Irritability. Loss of appetite. Malaise. 	 Fever. Mild rash (not infectious). Swelling of glands in the cheeks or neck. If these side effects occur, it is usually within 6-14 days of the vaccination.
Meningitis B (Men B)	,
 Pain, redness and swelling at the injection site. Fever. Irritability. Diarrhoea. Crying. 	by Bénédiere Barrett This Photo by Unknown Author is licensed under CC.

What to do:

- Give extra fluids.
- Do not overdress baby/child.
- Give paracetamol if needed for pain or to lower temperature.

*If adverse events following immunisation are severe and persistent or you have any concerns about your child seek medical advice either by contacting the surgery, calling 111 or 999 or visiting www.what0-18.nhs.uk